## **AMENDMENTS TO THE CLAIMS**

Docket No.: 13477-00002-US

## **Listing of claims:**

- 1. (Original) A method for generating or increasing the resistance of a plant to a plant pathogen of the phylum Oomyceta comprising increasing the activity of Rpi-blb2 protein in the plant or a tissue, organ or cell of a plant or a part thereof.
- 2. (Currently amended) The method of claim 1, wherein said Rpi-blb2 protein is encoded by a polynucleotide comprising a nucleic acid molecule selected from the group consisting of:
- (a) nucleic acid molecule encoding at least the mature form of the polypeptide depicted in SEQ ID NO: 2 or 4;
- (b) nucleic acid molecule comprising the coding sequence as depicted in SEQ ID NO: 1 or 3 or 5 or 6 encoding at least the mature form of the polypeptide;
- (c) nucleic acid molecules the nucleotide sequence of which is degenerate as a result of the genetic code to a nucleotide sequence of (a) or (b);
- (d) nucleic acid molecule encoding a polypeptide derived from the polypeptide encoded by a polynucleotide of (a) to (c) by way of substitution, deletion and/or addition of one or several amino acids of the amino acid sequence of the polypeptide encoded by a polynucleotide of (a) to (c);
- (e) nucleic acid molecule encoding a polypeptide the sequence of which has an identity of 70% or more to the amino acid sequence of the polypeptide encoded by a nucleic acid molecule of (a) or (b);
- (f) nucleic acid molecule comprising a fragment or a epitope-bearing portion of a polypeptide encoded by a nucleic acid molecule of any one of (a) to (e);
- (g) nucleic acid molecule comprising a polynucleotide having a sequence of a nucleic acid molecule amplified from a nucleic acid library using a primer as listed in Tab. 3b;

polypeptide encoded by any one of (a) to (g);

(h) nucleic acid molecule encoding a fragment beginning with amino acid: 1, 30, 50, 100, 200, 300, 500, or 1000 and stopping with amino acid 1276, 1000, 500, 300, 200, 50, or 1 of a

Docket No.: 13477-00002-US

- (i) nucleic acid molecule comprising at least 20 nucleotides of a polynucleotide of any one of (a) or (d);
- (j) nucleic acid molecule encoding a polypeptide being recognized by a monoclonal antibody that have been raised against a polypeptide encoded by a nucleic acid molecule of any one of (a) to (h);
- (k) nucleic acid molecule obtainable by screening an appropriate library under stringent conditions with a probe having the sequence of the nucleic acid molecule of any one of (a) to (j) or of a fragment thereof of at least 20; and
- (l) nucleic acid molecule the complementary strand of which hybridises under stringent conditions with a nucleic acid molecule of any one of (a) or (k);

or the complementary strand of any one of (a) to (1);

or expressing a polypeptide encoded by a segment of chromosome or linkage group 6 of Solanum bulbocastanum or Solanum tuberosum which co-segregates with a marker selected from table 3a or 3b and which mediates resistance to a pathogen of the phylum Oomyceta

and whereby the polynucleotide has not does not have the sequence of Mi1.1 or Mi1.2 as depicted in Seq. ID NO: SEQ ID NO: 7 or 9.

- 3. (Currently amended) The method of claim 1 [or 2], wherein the activity of a further resistance protein is increased.
- 4. (Currently amended) The method of any one of claims 1-to 3 claim 1, wherein activity is increased due to a de novo-expression.

- 5. (Currently amended) The method of any one of claims 1 to 4 claim 1, wherein the endogenous activity of a Rpi-blb2 and/or the further resistance protein is increased.
- 6. (Currently amended) The method of any one of claim 1 to 5 claim 1, comprising one or more of the following steps

- a) stabilizing the resistance protein;
- b) stabilizing the resistance protein encoding mRNA;
- c) increasing the specific activity of the resistance protein;
- d) expressing or increasing the expression of a homologous or artificial transcription factor for resistance protein expression;
- e) stimulate resistance protein activity through exogenous inducing factors;
- f) expressing a transgenic resistance protein encoding gene; and/or
- g) increasing the copy number of the resistance protein encoding gene.
- 7. (Currently amended) The method of any one of claims 1 to 6 claim 1 which results in reduction in the sporulation index of at least 30% after infection with P. infestans compared to a wild type.
- 8. (Currently amended) A polynucleotide encoding a Rpi-blb2 protein comprising a nucleic acid molecule selected from the group consisting of:
- (a) nucleic acid molecule encoding at least the mature form of the polypeptide depicted in SEQ ID NO: 2 or 4;
- (b) nucleic acid molecules comprising the coding sequence as depicted in SEQ ID NO: 1 or 3 or 5 or 6 encoding at least the mature form of the polypeptide;
- (c) nucleic acid molecule the nucleotide sequence of which is degenerate as a result of the genetic code to a nucleotide sequence of (a) or (b);

(d) nucleic acid molecule encoding a polypeptide derived from the polypeptide encoded by a polynucleotide of (a) to (c) by way of substitution, deletion and/or addition of one or several amino acids of the amino acid sequence of the polypeptide encoded by a polynucleotide of (a) to (c);

Docket No.: 13477-00002-US

- (e) nucleic acid molecule encoding a polypeptide the sequence of which has an identity of 70% or more to the amino acid sequence of the polypeptide encoded by a nucleic acid molecule of (a) or (b);
- (f) nucleic acid molecules comprising a fragment or a epitope-bearing portion of a polypeptide encoded by a nucleic acid molecule of any one of (a) to (e);
- (g) nucleic acid molecule comprising a polynucleotide having a sequence of a nucleic acid molecule amplified from a nucleic acid library using the primers as listed in Tab.3b;
- (h) nucleic acid molecule encoding polypeptide fragment beginning with amino acid: 1, 30, 50, 100, 200, 300, 500, or 1000 and stopping with amino acid 1276, 1000, 500, 300, 200, 50, or 30 of a polypeptide encoded by any one of (a) to (g);
- (i) nucleic acid molecule comprising at least 20 nucleotides of a polynucleotide of any one of (a) or (d);
- (j) nucleic acid molecule encoding a polypeptide being recognized by a monoclonal antibodies that have been raised against a polypeptide encoded by a nucleic acid molecule of any one of (a) to (h);
- (k) nucleic acid molecule obtainable by screening an appropriate library under stringent conditions with a probe having the sequence of the nucleic acid molecule of any one of (a) to (j) or of a fragment thereof of at least 20; and
- (l) nucleic acid molecule the complementary strand of which hybridises under stringent conditions with a nucleic acid molecule of any one of (a) or (k);

or the complementary strand of any one of (a) to (l);

mediates resistance to pathogens of the phylum Oomyceta;

or encoding a polypeptide encoded by a segment of chromosome or of linkage group 6 of Solanum bulbocastanum or Solanum tuberosum which co-segregates with a marker selected from table 3a or 3b or comprises a replication site or hybridisation site for said marker and which

Docket No.: 13477-00002-US

and whereby the polynucleotide does not consist of the sequence shown in Rossi et al. 1998, PNAS USA 95:9750-9754, Milligan et al., 1998, Plant Cell 10: 1307-1319, or WO 98/06750 of Mil.1 or Mil.2 as depicted in SEQ ID NO: 7 or 9.

- 9. (Currently amended) The polynucleotide of claim 8 or the method of any one of claims 1 to 7, wherein the marker is E40M58, CT119, or CT216.
- 10. (Currently amended) The polynucleotide of claim 8 [to 9] which is DNA or RNA.
- 11. (Currently amended) A method for making a recombinant vector comprising inserting the polynucleotide of any one of claims 8 to 10 claim 8 into a vector or inserting said polynucleotide and a further resistance protein.
- 12. (Currently amended) A vector containing the polynucleotide of any one of claims 8 to 10 claim 8 or comprising said polynucleotide and a further resistance gene or being produced by the method of claim 11.
- 13. (Currently amended) The vector of claim 12 or the method of any one of claims 1 to 7 in which a polynucleotide encoding Rpi-blb2 protein or encoding the further resistance protein is operatively linked to expression control sequences and/or is operatively linked to a nucleic acid sequence encoding a transgenic expression regulating signal allowing expression in prokaryotic or eukaryotic host cells.
- 14. (Currently amended) The vector of claim 12 [or 13] or the method of any one of claims 1 to 7 in which the polynucleotide encoding Rpi-blb2 protein or encoding a further resistance protein is operatively linked to expression control sequences of the same species origin as the polynucleotide encoding Rpi-blb2 protein or the further resistance protein.

15. (Currently amended) A method of making a recombinant host cell comprising introducing the vector of any one of claims 12 to 14 claim 12 or introducing said vector and a vector for expressing a further resistance protein into a host cell.

- 16. (Currently amended) A host cell produced according to the method of claim 15 of genetically engineered with the polynucleotide of any one of claims 8 to 10 or the vector of any one of claims 12 to 14 or genetically engineered with said vector or polynucleotide and a vector or a polynucleotide for expressing a further resistance protein.
- 17. (Original) The host cell of claim 16, which is E. coli, Baculovirus, Agrobacterium, or a plant cell.
- 18. (Currently amended) A process for the production of a Rpi-blb2-polypeptide comprising culturing the host cell of claim 16 [or 17] and recovering the polypeptide encoded by said polynucleotide and expressed by the host cell from the culture or the host cells.
- 19. (Currently amended) A polypeptide having the amino acid sequence encoded by a polynucleotide of any one of claims 8 to 10 claim 8 or obtainable by the process of claim 18.
- 20. (Original) A polypeptide having Rpi-blb2 activity.
- 21. (Currently amended) An antibody that binds specifically to the polypeptide of claim 19 [or 20].
- 22. (Currently amended) An antisense nucleic acid molecule comprising the complementary sequence of the polynucleotide of any-one of claims 8 to 10 claim 8.
- 23. (Currently amended) A method for the production of a transgenic plant, plant cell or plant tissue or a part thereof comprising the introduction of introducing the polynucleotide of any one of claims 8 to 10 claim 8 or said polynucleotide and a polynucleotide encoding a further resistance protein, or the vector of any one of claims 12 to 14 into the genome of said plant, plant tissue or plant cell or a part thereof.

24. (Currently amended) A plant cell comprising the polynucleotide of any one of claims 8 to 10 claim 8, the vector of any one of claims 12 to 14 or obtainable by the method of claim 23 or a vector containing the polynucleotide of claim 8 or comprising the polynucleotide of claim 8 and a further resistance gene.

Docket No.: 13477-00002-US

- 25. (Original) A transgenic plant or plant tissue or a part thereof comprising the plant cell of claim 24.
- 26. (Currently amended) A method for producing a plant or a part thereof resistant to a plant pathogen of the phylum Oomyceta comprising the step:

expressing in the plant or a part thereof the polypeptide of claim 19 [or 20] and a further resistance protein.

- 27. (Currently amended) A method for producing a plant or a part thereof with a durable resistance to a Phytophthora sp. comprising co-expressing in the plant or a part thereof Rpi-blb and Rpi-blb2 protein or the polypeptide of claim 19 [or 20].
- 28. (Currently amended) The transgenic plant or plant tissue of claim 25 or produced according to claim 26 or 27, which upon the presence of the polynucleotide or the vector is resistant to a plant pathogen of the phylum Oomyceta.
- 29. (Currently amended) Harvestable parts of the transgenic plant or plant tissue of claim 25 comprising the plant cell of claim 24.
- 30. (Currently amended) Propagation material of the transgenic plant or plant tissue of claim 25 comprising the plant cell of claim 24.
- 31. (Currently amended) Use of the polynucleotide of any one of claims 8 to 10, the vector of any one of claims 12 to 14, or the polypeptide of claim 19 or 20 A method for producing a plant or a plant tissue, plant organ, or a plant cell or a part thereof resistant to a plant pathogen of the phylum Oomyceta comprising introducing the vector of claim 12 into a plant, plant tissue or plant cell or a part thereof.

32. (Currently amended) A method for the identification of an a compound stimulating resistance to a plant pathogen of the phylum Oomyceta comprising:

- (a) contacting cells which express the polypeptide of claim 19 [or 20] or its mRNA with a candidate compound under cell cultivation conditions;
- (b) assaying an increase in expression of said polypeptide or said mRNA;
- (c) comparing the expression level to a standard response made in the absence of said candidate compound; whereby, an increased expression over the standard indicates that the compound is stimulating resistance.
- 33. (Canceled)
- 34. (Currently amended) A diagnostic composition, comprising the polynucleotide of any one of claims 8 to 10 claim 8, the a vector—of any one of claims 12 to 14, the an antibody that binds specifically to a polypeptide having the amino acid sequence encoded by the polynucleotide of claim 8 of claim 21 or the or an antisense nucleic acid of claim 22 molecule comprising the complementary sequence of the polynucleotide of claim 8 and optionally suitable means for detection.
- 35. (Currently amended) A kit comprising the polynucleotide of any one of claims 8 or 12 claim 8, the a vector of any one of claims 12 to 14, the host cell of claim 16 or 17, the polypeptide of claim 19 or 20, the antisense nucleic acid of claim 22, the antibody of claim 21, the plant cell of claim 24, the plant or plant tissue of claim 25 comprising the polynucleotide of claim 8, the or harvestable part of claim 29, or the propagation material derived therefrom, of claim 30 or an antibody that binds specifically to a polypeptide having the amino acid sequence encoded by the polynucleotide of claim 8 or an antisense nucleic acid molecule comprising the complementary sequence of the polynucleotide of claim 8 and optionally a polynucleotide encoding Rpi blb, a Rpi blb protein or an antibody against Rpi blb.
- 36. (Currently amended) A method for the production of a plant crop protectant <u>comprising</u> providing the polynucleotide of any one of claims 8 to 10 claim 8, the vector of any one of

claims 12 to 14 or the polypeptide of claim 19 or 20 or comprising the steps of the method of claim 32; and formulating the polynucleotide of any one of claims 8 to 10, the vector of of claims 12 or 14 or the polypeptide of claim 19 or 20 or the compound identified in step (c) of claim 32 in a form applicable as an agricultural composition.

- 37. (Currently amended) The <u>polynucleotide of claim 8 or a vector</u>, host cell, plant tissue, plant, <u>use</u>, <u>or</u> kit <del>or method of any one of claims 1 to 36</del> <u>comprising the polynucleotide of claim 8</u>, wherein the plant pathogen is of the order Pythiales or Peronosperales.
- 38. (Currently amended) The <u>polynucleotide of claim 8 or a vector</u>, host cell, plant cell, plant tissue, plant, use, or kit or method of any one of claims 1 to 37 comprising the polynucleotide of <u>claim 8</u>, wherein the <u>plant-pathogen</u> is of the species P. infestans, Phytophthora erythroseptica, Phytophthora capsici, Phytophthora sojae, Phytophthora parasitica var. nicotianae, Bremia lactuca, Peronospera tabaci or Plasmopara viticola.
- 39. (Currently amended) The vector, host cell, plant cell, plant tissue, plant, use, kit or method of any one of claims 1 to 38 claim 1, wherein the resistance protein is characterized by a P-loop and a NBS domain.
- 40. (Currently amended) The vector of claim 12, or a host cell, plant cell, plant tissue, plant, use, or kit or method of any one of claims 1 to 39 comprising the vector of claim 12, wherein the further resistance gene is a gene encoding Rpi-blb, R1, R-ber, Rpi1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R11, Ph-1, Ph-2 and/or Ph-3.
- 41. (Currently amended) The vector of claim 12, or a host cell, plant cell, plant tissue, plant, use, or kit or method of any one of claims 1 to 40 comprising the vector of claim 12, wherein the further resistance protein is the Rpi-blb protein.
- 42. (Currently amended) The vector, host cell, plant cell of claim 24, or plant tissue, plant, use, or kit or method of any one of claims 1 to 41 comprising the plant cell of claim 24 wherein the plant, plant cell or plant tissue is selected from the group consisting of Menyanthaceae, Solanaceae, Sclerophylacaceae, Duckeodendraceae, Goetzeaceae, Convolvulaceae, Cuscutaceae,

Polemoniaceae, and Hydrophyllaceae according to the Systema Naturae 2000, Brands, S.J., Amsterdam or has its origin thereof.

Docket No.: 13477-00002-US

43. (Currently amended) The vector, polynucleotide of claim 8, a polypeptide encoded by the polynucleotide of claim 8, a host cell, plant cell, plant tissue, or plant comprising the polynucleotide of claim 8, use, kit or method of any one of claims 1 to 42, wherein the polynucleotide, the polypeptide, the plant cell, the host cell, the plant tissue or the plant is derived from a member of the Solanceae Solanaceae family, preferably selected from the group consisting of S. bulbocastanum, potato (S. tuberosum), tomato (S. lycopersicum or Lycopersicum lycopersicum (L.) Karsten ex Farwell), petunia, tree tomato (S. betaceum), pear melon (S. muricatum) of and eggplant (S. melongena).